

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Application No. :

U.S. National Serial No. :

Filed :

PCT International Application No. : PCT/DE00/00244

VERIFICATION OF A TRANSLATION

I, Susan POTTS BA ACIS

Director to RWS Group plc, of Europa House, Marsham Way, Gerrards Cross, Buckinghamshire, England declare:

That the translator responsible for the attached translation is knowledgeable in the German language in which the below identified international application was filed, and that, to the best of RWS Group plc knowledge and belief, the English translation of the international application No. PCT/DE00/00244 is a true and complete translation of the amended claims of the PCT of the above identified international application as filed.

I hereby declare that all the statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the patent application issued thereon.

Date: June 15, 2001

Signature of Director :



For and on behalf of RWS Group plc

Post Office Address :

Europa House, Marsham Way,  
Gerrards Cross, Buckinghamshire,  
England.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Application No. :

U.S. National Serial No. :

Filed :

PCT International Application No. : PCT/DE00/00244

VERIFICATION OF A TRANSLATION

I, the below named translator, hereby declare that:

My name and post office address are as stated below;

That I am knowledgeable in the German language in which the below identified international application was filed, and that, to the best of my knowledge and belief, the English translation of the international application No. PCT/DE00/00244 is a true and complete translation of the above identified international application as filed.

I hereby declare that all the statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the patent application issued thereon.

Date: July 10, 2001

Full name of the translator :

  
Margrit METHUEN

For and on behalf of RWS Group plc

Post Office Address :

Europa House, Marsham Way,  
Gerrards Cross, Buckinghamshire,  
England.

5 / PRTS

889802  
JC17 Re PCT/PTO 20 JUL 2001

WO 00/44895

PCT/DE00/00244

**Method and medicament for inhibiting the expression of a given gene**

E'   
 5   
 SUB   
 E2

The invention relates to methods in accordance with the preambles of claims 1 and 2. It furthermore relates to a medicament and to a use of double-stranded oligoribonucleotides and to a vector encoding them.

10 Such a method is known from WO 99/32619, which was unpublished at the priority date of the present invention. The known process aims at inhibiting the expression of genes in cells of invertebrates. To this end, the double-stranded oligoribonucleotide must exhibit a sequence which is identical with the target

15 gene and which has a length of at least 50 bases. To achieve efficient inhibition, the identical sequence must be 300 to 1 000 base pairs in length. Such an oligoribonucleotide is complicated to prepare.

20 DE 196 31 919 C2 describes an antisense RNA with specific secondary structures, the antisense RNA being present in the form of a vector encoding it. The antisense RNA takes the form of an RNA molecule which is complementary to regions of the mRNA. Inhibition of

25 the gene expression is caused by binding to these regions. This inhibition can be employed in particular for the diagnosis and/or therapy of diseases, for example tumor diseases or viral infections. - The disadvantage is that the antisense RNA must be

30 introduced into the cell in an amount which is at least as high as the amount of the mRNA. The known antisense methods are not particularly effective.

US 5,712,257 discloses a medicament comprising

35 mismatched double-stranded RNA (dsRNA) and bioactive mismatched fragments of dsRNA in the form of a ternary complex together with a surfactant. The dsRNA used for this purpose consists of synthetic nucleic acid single strands without defined base sequence. The single

strands undergo irregular base pairing, also known as "non-Watson-Crick" base pairing, giving rise to mismatched double strands. The known dsRNA is used to inhibit the amplification of retroviruses such as HIV.

5 Amplification of the virus can be inhibited when non-sequence-specific dsRNA is introduced into the cells. This leads to the induction of interferon, which is intended to inhibit viral amplification. The inhibitory effect, or the activity, of this method is poor.

10

It is known from Fire, A. et al., NATURE, Vol. 391, pp. 806 that dsRNA whose one strand is complementary in segments to a nematode gene to be inhibited inhibits the expression of this gene highly efficiently. It is  
15 believed that the particular activity of the dsRNA used in nematode cells is not due to the antisense principle but possibly on catalytic properties of the dsRNA, or enzymes induced by it. - Nothing is mentioned in this paper on the activity of specific dsRNA with regard to  
20 inhibiting the gene expression, in particular in mammalian and human cells.

The object of the present invention is to do away with the disadvantages of the prior art. In particular, it  
25 is intended to provide as effective as possible a method, medicament or use for the preparation of a medicament, which method, medicament or use is capable of causing particularly effective inhibition of the expression of a given target gene.

30

This object is achieved by the features of claims 1, 2, 37, 38 and 74 and 75. Advantageous embodiments can be seen from claims 3 to 36, 39 to 73 and 76 to 112.

*SUB E3*  
*E4* 35 In accordance with the method-oriented inventions, it is provided in each case that the region I which is complementary to the target gene exhibits not more than 49 successive nucleotide pairs.

Provided in accordance with the invention are an oligoribonucleotide or a vector encoding therefor. At least segments of the oligoribonucleotide exhibit a defined nucleotide sequence. The defined segment may be  
5 limited to the complementary region I. However, it is also possible that all of the double-stranded oligoribonucleotide exhibits a defined nucleotide sequence.

10 Surprisingly, it has emerged that an effective inhibition of the expression of the target gene can be achieved even when the complementary region I is not more than 49 base pairs in length. The procedure of providing such oligoribonucleotides is less  
15 complicated.

In particular, dsRNA with a length of over 50 nucleotide pairs induces certain cellular mechanisms, for example the dsRNA-dependent protein  
20 kinase or the 2-5A system, in mammalian and human cells. This leads to the disappearance of the interference effect mediated by the dsRNA which exhibits a defined sequence. As a consequence, protein biosynthesis in the cell is blocked. The present  
25 invention overcomes this disadvantage in particular.

Furthermore, the uptake of dsRNA with short chain lengths into the cell or into the nucleus is facilitated markedly over longer-chain dsRNAs.

30 It has proved advantageous for the dsRNA or the vector to be present packaged into micellar structures, preferably in liposomes. The dsRNA or the vector can likewise be enclosed in viral natural capsids or in  
35 chemically or enzymatically produced artificial capsids or structures derived therefrom. - The abovementioned features make it possible to introduce the dsRNA or the vector into given target cells.

In a further aspect, the dsRNA has 10 to 1 000, preferably 15 to 49, base pairs. Thus, the dsRNA can be longer than the region I, which is complementary to the target gene. The complementary region I can be located  
5 at the terminus or inserted into the dsRNA. Such dsRNA or a vector provided for coding the same can be produced synthetically or enzymatically by customary methods.

10 The gene to be inhibited is expediently expressed in eukaryotic cells. The target gene can be selected from the following group: oncogene, cytokin gene, Id protein gene, developmental gene, prion gene. It can also be expressed in pathogenic organisms, preferably in  
15 plasmodia. It can be part of a virus or viroid which is preferably pathogenic to humans. - The method proposed makes it possible to produce compositions for the therapy of genetically determined diseases, for example cancer, viral diseases or Alzheimer's disease.

20 The virus or viroid can also be a virus or viroid which is pathogenic to animals or plant-pathogenic. In this case, the method according to the invention also permits the provision of compositions for treating  
25 animal or plant diseases.

In a further aspect, segments of the dsRNA are designed as double-stranded. A region II which is complementary within the double-stranded structure is formed by two  
30 separate RNA single strands or by autocomplementary regions of a topologically closed RNA single strand which is preferably in circular form.

The ends of the dsRNA can be modified to counteract  
35 degradation in the cell or dissociation into the single strands. Dissociation takes place in particular when low concentrations or short chain lengths are used. To inhibit dissociation in a particularly effective fashion, the cohesion of the complementary region II,

which is caused by the nucleotide pairs, can be increased by at least one, preferably two, further chemical linkage(s). - A dsRNA according to the invention whose dissociation is reduced exhibits  
5 greater stability to enzymatic and chemical degradation in the cell or in the organism.

The complementary region II can be formed by autocomplementary regions of an RNA hairpin loop, in  
10 particular when using a vector according to the invention. To afford protection from degradation, it is expedient for the nucleotides to be chemically modified in the loop region between the double-stranded structure.

15 The chemical linkage is expediently formed by a covalent or ionic bond, a hydrogen bond, hydrophobic interactions, preferably van-der-Waals or stacking interactions, or by metal-ion coordination. In an  
20 especially advantageous aspect, it can be formed at at least one, preferably both, end(s) of the complementary region II.

It has furthermore proved to be advantageous for the  
25 chemical linkage to be formed by one or more linkage groups, the linkage groups preferably being poly(oxyphosphinicoxy-1,3-propanediol) and/or polyethylene glycol chains. The chemical linkage can also be formed by purine analogs used in place of purines in  
30 the complementary regions II. It is also advantageous for the chemical linkage to be formed by azabenzene units introduced into the complementary regions II. Moreover, it can be formed by branched nucleotide analogs used in place of nucleotides in the  
35 complementary regions II.

It has proved expedient to use at least one of the following groups for generating the chemical linkage: methylene blue; bifunctional groups, preferably

bis(2-chloroethyl)amine; N-acetyl-N'-(p-glyoxyl-  
benzoyl)cystamine; 4-thiouracil; psoralene. The  
chemical linkage can furthermore be formed by  
thiophosphoryl groups provided at the ends of the  
5 double-stranded region. The chemical linkage at the  
ends of the double-stranded region is preferably formed  
by triple-helix bonds.

10 The chemical linkage can expediently be induced by  
ultraviolet light.

The nucleotides of the dsRNA can be modified. This  
counteracts the activation, in the cell, of a double-  
stranded-RNA-dependent protein kinase, PKR.  
15 Advantageously, at least one 2'-hydroxyl group of the  
nucleotides of the dsRNA in the complementary region II  
is replaced by a chemical group, preferably a 2'-amino  
or a 2'-methyl group. At least one nucleotide in at  
least one strand of the complementary region II can  
20 also be a locked nucleotide with a sugar ring which is  
chemically modified, preferably by a 2'-O, 4'-C  
methylene bridge. Advantageously, several nucleotides  
are locked nucleotides.

25 A further especially advantageous embodiment provides  
that the dsRNA or the vector is bound to, associated  
with or surrounded by, at least one viral coat protein  
which originates from a virus, is derived therefrom or  
has been prepared synthetically. The coat protein can  
30 be derived from polyomavirus. The coat protein can  
contain the polyomavirus virus protein 1 (VP1) and/or  
virus protein 2 (VP2). The use of such coat proteins is  
known from, for example, DE 196 18 797 A1, whose  
disclosure is herewith incorporated. - The  
35 abovementioned features considerably facilitate the  
introduction of the dsRNA or of the vector into the  
cell.



When a capsid or capsid-type structure is formed from the coat protein, one side preferably faces the interior of the capsid or capsid-type structure. The construct formed is particularly stable.

5

The dsRNA can be complementary to the primary or processed RNA transcript of the target gene. - The cell can be a vertebrate cell or a human cell.

- 10 At least two dsRNAs which differ from each other or at least one vector encoding them can be introduced into the cell, where at least segments of one strand of each dsRNA are complementary to in each case one of at least two different target genes. This makes it possible  
15 simultaneously to inhibit the expression of at least two different target genes. In order to suppress, in the cell, the expression of a double-stranded-RNA-dependent protein kinase, PKR, one of the target genes is advantageously the PKR gene. This allows effective  
20 suppression of the PKR activity in the cell.

- The invention furthermore provides a medicament with at least one oligoribonucleotide with double-stranded structure (dsRNA) for inhibiting the expression of a  
25 given target gene, where one strand of the dsRNA has a region I where at least segments are complementary to the target gene. - Surprisingly, it has emerged that such a dsRNA is suitable as medicament for inhibiting the expression of a given gene in mammalian cells. In  
30 comparison with the use of single-stranded oligoribonucleotides, the inhibition is already caused at concentrations which are lower by at least one order of magnitude. The medicament according to the invention is highly effective. Lesser side effects can be  
35 expected.

The invention furthermore provides a medicament with at least one vector for coding at least one oligoribonucleotide with double-stranded structure

(dsRNA) for inhibiting the expression of a given target gene, where one strand of the dsRNA has a region I where at least segments are complementary to the target gene. - The medicament proposed exhibits the  
5 abovementioned advantages. By using a vector, in particular production costs can be reduced.

In a particularly advantageous embodiment, the complementary region I has not more than 49 successive  
10 nucleotide pairs. - Surprisingly, it has emerged that an effective inhibition of the expression of the target gene can be achieved even when the complementary region I is not more than 49 base pairs in length. The procedure of providing such oligoribonucleotides is  
15 less complicated.

The invention furthermore provides a use of an oligoribonucleotide with double-stranded structure (dsRNA) for preparing a medicament for inhibiting the  
20 expression of a given target gene, where one strand of the dsRNA has a region I where at least segments are complementary to the target gene. - Surprisingly, such a dsRNA is suitable for preparing a medicament for inhibiting the expression of a given gene. Compared  
25 with the use of single-stranded oligoribonucleotides, the inhibition is already caused at concentrations which are lower by one order of magnitude when using dsRNA. The use according to the invention thus makes possible the preparation of particularly effective  
30 medicaments.

The invention furthermore provides the use of a vector for coding at least one oligoribonucleotide with double-stranded structure (dsRNA) for preparing a  
35 medicament for inhibiting the expression of a given target gene, where one strand of the dsRNA has a region I where at least segments are complementary to this target gene. - The use of a vector makes possible a particularly effective gene therapy.

With regard to advantageous embodiments of the medicament and of the use, reference is made to the description of the above features.

5

Use examples of the invention are illustrated in greater detail hereinbelow with reference to the figures, in which:

10 Fig. 1 shows the schematic representation of a plasmid for the *in vitro* transcription with T7- and SP6-polymerase,

15 Fig. 2 shows RNA following electrophoresis on an 8% polyacrylamide gel and staining with ethidium bromide,

20 Fig. 3 shows a representation of radioactive RNA transcripts following electrophoresis on an 8% polyacrylamide gel with 7 M urea by means of an instant imager, and

Figs. 4a - e show Texas Red and YFP fluorescence in murine fibroblasts.

25

**Use example 1:**

The inhibition of transcription was detected by means of sequence homologous dsRNA in an *in vitro* transcription system with a nuclear extract from human  
30 HeLa cells. The DNA template for this experiment was plasmid pCMV1200 which had been linearized by means of *Bam*HI.

**Generation of the template plasmids:**

35 The plasmid shown in fig. 1 was constructed for use in the enzymatic synthesis of the dsRNA. To this end, a polymerase chain reaction (PCR) with the "positive control DNA" of the HeLaScribe® Nuclear Extract *in vitro* transcription kit by Promega, Madison, USA, as

DNA template was first carried out. One of the primers used contained the sequence of an *EcoRI* cleavage site and of the T7 RNA polymerase promoter as shown in sequence listing No. 1. The other primer contained the sequence of a *BamHI* cleavage site and of the SP6 RNA polymerase promoter as shown in sequence listing No. 2. In addition, the two primers had, at the 3' ends, regions which were identical with or complementary to the DNA template. The PCR was carried out by means of the "Taq PCR Core Kits" by Qiagen, Hilden, Germany, following the manufacturer's instructions. 1.5 mM  $MgCl_2$ , in each case 200  $\mu M$  dNTP, in each case 0.5  $\mu M$  primer, 2.5 U Taq DNA polymerase and approximately 100 ng of "positive control DNA" were employed as template in PCR buffer in a volume of 100  $\mu l$ . After initial denaturation of the template DNA by heating for 5 minutes at 94°C, amplification was carried out in 30 cycles of denaturation for in each case 60 seconds at 94°C, annealing for 60 seconds at 5°C below the calculated melting point of the primers and polymerization for 1.5-2 minutes at 72°C. After a final polymerization of 5 minutes at 72°C, 5  $\mu l$  of the reaction were analyzed by agarose-gel electrophoresis. The length of the DNA fragment amplified thus was 400 base pairs, 340 base pairs corresponding to the "positive control DNA". The PCR product was purified, hydrolyzed with *EcoRI* and *BamHI* and, after repurification, employed in the ligation together with a pUC18 vector which had also been hydrolyzed by *EcoRI* and *BamHI*. *E. coli* XL1-blue was then transformed. The plasmid obtained (pCMV5) carries a DNA fragment whose 5' end is flanked by the T7 promoter and whose 3' end is flanked by the SP6 promoter. By linearizing the plasmid with *BamHI*, it can be employed *in vitro* with the T7-RNA polymerase for the run-off transcription of a single-stranded RNA which is 340 nucleotides in length and shown in sequence listing No. 3. If the plasmid is linearized with *EcoRI*, it can be employed for the run-off transcription with SP6 RNA polymerase,

giving rise to the complementary strand. In accordance with the method outlined hereinabove, an RNA 23 nucleotides in length was also synthesized. To this end, a DNA shown in sequence listing No. 4 was ligated with the pUC18 vector via the *EcoRI* and *BamHI* cleavage sites.

Plasmid pCMV1200 was constructed as DNA template for the *in-vitro* transcription with HeLa nuclear extract. To this end, a 1 191 bp *EcoRI*/*BamHI* fragment of the positive control DNA contained in the HeLaScribe® Nuclear Extract *in vitro* transcription kit was amplified by means of PCR. The amplified fragment encompasses the 828 bp "immediate early" CMV promoter and a 363 bp transcribable DNA fragment. The PCR product was ligated to the vector pGEM-T via "T-overhang" ligation. A *BamHI* cleavage site is located at the 5' end of the fragment. The plasmid was linearized by hydrolysis with *BamHI* and used as template in the run-off transcription.

*In-vitro* transcription of the complementary single strands:

pCMV5 plasmid DNA was linearized with *EcoRI* or *BamHI*. It was used as DNA template for an *in-vitro* transcription of the complementary RNA single strands with SP6 and T7 RNA polymerase, respectively. The "Riboprobe *in vitro* Transcription" system by Promega, Madison, USA, was employed for this purpose. Following the manufacturer's instructions, 2 µg of linearized plasmid DNA were incubated in 100 µl of transcription buffer and 40 U T7 or SP6 RNA polymerase for 5-6 hours at 37°C. The DNA template was subsequently degraded by addition of 2.5 µl of RNase-free DNase RQ1 and incubation for 30 minutes at 37°C. The transcription reaction was made up to 300 µl with H<sub>2</sub>O and purified by phenol extraction. The RNA was precipitated by addition of 150 µl of 7 M ammonium acetate [sic] and 1 125 µl of

ethanol and stored at  $-65^{\circ}\text{C}$  until used for the hybridization.

Generation of the RNA double strands:

- 5 For the hybridization, 500  $\mu\text{l}$  of the single-stranded RNA which had been stored in ethanol and precipitated were spun down. The resulting pellet was dried and taken up in 30  $\mu\text{l}$  of PIPES buffer, pH 6.4 in the presence of 80% formamide, 400 mM NaCl and 1 mM EDTA.
- 10 In each case 15  $\mu\text{l}$  of the complementary single strands were combined and heated for 10 minutes at  $85^{\circ}\text{C}$ . The reactions were subsequently incubated overnight at  $50^{\circ}\text{C}$  and cooled to room temperature.
- 15 Only approximately equimolar amounts of the two single strands were employed in the hybridization. This is why the dsRNA preparations contained single-stranded RNA (ssRNA) as contaminant. In order to remove these ssRNA contaminants, the reactions were treated, after
- 20 hybridization, with the single-strand-specific ribonucleases bovine pancreatic RNase A and *Aspergillus oryzae* RNase T1. RNase A is an endoribonuclease which is specific for pyrimidines. RNase T1 is an endoribonuclease which preferentially cleaves at the 3'
- 25 side of guanosines. dsRNA is no substrate for these ribonucleases. For the RNase treatment, the reactions in 300  $\mu\text{l}$  of Tris, pH 7.4, 300 mM NaCl and 5 mM EDTA were treated with 1.2  $\mu\text{l}$  of RNaseA at a concentration of 10 mg/ml and 2  $\mu\text{l}$  of RNaseT1 at a concentration of
- 30 290  $\mu\text{g/ml}$ . The reactions were incubated for 1.5 hours at  $30^{\circ}\text{C}$ . Thereupon, the RNases were denatured by addition of 5  $\mu\text{l}$  of proteinase K at a concentration of 20 mg/ml and 10  $\mu\text{l}$  of 20% SDS and incubation for 30 minutes at  $37^{\circ}\text{C}$ . The dsRNA was purified by phenol
- 35 extraction and precipitated with ethanol. To verify the completeness of the RNase digestion, two control reactions were treated with ssRNA analogously to the hybridization reactions.

The dried pellet was taken up in 15  $\mu$ l of TE buffer, pH 6.5, and subjected to native polyacrylamide gel electrophoresis on an 8% gel. The acrylamide gel was subsequently stained in an ethidium bromide solution and washed in a water bath. Fig. 2 shows the RNA which had been visualized in a UV transilluminator. The sense RNA which had been applied to lane 1 and the antisense RNA which had been applied to lane 2 showed a different migration behavior under the chosen conditions than the dsRNA of the hybridization reaction which had been applied to lane 3. The RNase-treated sense RNA and antisense RNA which had been applied to lanes 4 and 5, respectively, produced no visible band. This shows that the single-stranded RNAs had been degraded completely. The RNase-treated dsRNA of the hybridization reaction which had been applied to lane 6 is resistant to RNase treatment. The band which migrates faster in the native gel in comparison with the dsRNA applied to lane 3 results from dsRNA which is free from ssRNA. In addition to the dominant main band, weaker bands which migrate faster are observed after the RNase treatment.

In-vitro transcription test with human nuclear extract:

Using the HeLaScribe<sup>®</sup> Nuclear Extract in vitro transcription kit by Promega, Madison, USA, the transcription efficiency of the abovementioned DNA fragment which is present in plasmid pCMV1200 and homologous to the "positive control DNA" was determined in the presence of the dsRNA (dsRNA-CMV5) with sequence homology. Also, the effect of the dsRNA without sequence homology, which corresponds to the yellow fluorescent protein (YFP) gene (dsRNA-YRP), was studied. This dsRNA had been generated analogously to the dsRNA with sequence homology. The sequence of a strand of this dsRNA can be found in sequence listing No. 5. Plasmid pCMV1200 was used as template for the run-off transcription. It carries the "immediate early" cytomegalovirus promoter which is recognized by the eukaryotic RNA polymerase II, and a transcribable DNA

fragment. Transcription was carried out by means of the HeLa nuclear extract, which contains all the proteins which are necessary for transcription. By addition of [ $-\text{}^{32}\text{P}$ ]rGTP to the transcription reaction, radiolabeled

5 transcript was obtained. The [ $-\text{}^{32}\text{P}$ ]rGTP used had a specific activity of 400 Ci/mmol, 10 mCi/ml. 3 mM  $\text{MgCl}_2$ , in each case 400  $\mu\text{M}$  rATP, rCTP, rUTP, 16  $\mu\text{M}$  rGTP, 0.4  $\mu\text{M}$  [ $-\text{}^{32}\text{P}$ ]rGTP and depending on the experiment

10 1 fmol of linearized plasmid DNA and various amounts of dsRNA in transcription buffer were employed per reaction. Each batch was made up to a volume of 8.5  $\mu\text{l}$  with  $\text{H}_2\text{O}$ . The reactions were mixed carefully. To start the transcription, 4 U HeLa nuclear extract in a volume of 4  $\mu\text{l}$  were added and incubated for 60 minutes at

15 30°C. The reaction was stopped by addition of 87.5  $\mu\text{l}$  of quench mix which had been warmed to 30°C. To remove the proteins, the reactions were treated with 100  $\mu\text{l}$  of phenol/chloroform/isoamyl alcohol (25:24:1 v/v/v) saturated with TE buffer, pH 5.0, and the reactions

20 were mixed vigorously for 1 minute. For phase separation, the reactions were spun for approximately 1 minute at 12 000 rpm and the top phase was transferred into a fresh reaction vessel. Each reaction was treated with 250  $\mu\text{l}$  of ethanol. The reactions were

25 mixed thoroughly and incubated for at least 15 minutes on dry ice/methanol. To precipitate the RNA, the reactions were spun for 20 minutes at 12 000 rpm and 40°C. The supernatant was discarded. The pellet was dried in vacuo for 15 minutes and resuspended in 10  $\mu\text{l}$

30 of  $\text{H}_2\text{O}$ . Each reaction was treated with 10  $\mu\text{l}$  of denaturing loading buffer. The free GTP was separated from the transcript formed by means of denaturing polyacrylamide gel electrophoresis on an 8% gel with 7 M urea. The RNA transcripts formed upon transcription

35 with HeLa nuclear extract, in denaturing loading buffer, were heated for 10 minutes at 90°C and 10  $\mu\text{l}$  aliquots were applied immediately to the freshly washed pockets. The electrophoresis was run at 40 mA. The amount of the radioactive ssRNA formed upon



transcription was analyzed after electrophoresis with the aid of an *Instant Imager*.

Fig. 3 shows the radioactive RNA from a representative test, shown by means of the *Instant Imager*. Samples obtained from the following transcription reactions were applied:

- Lane 1: without template DNA, without dsRNA;
- 10 Lane 2: 50 ng of template DNA, without dsRNA;
- Lane 3: 50 ng of template DNA, 0.5  $\mu$ g of dsRNA YFP;
- Lane 4: 50 ng of template DNA, 1.5  $\mu$ g of dsRNA YFP;
- Lane 5: 50 ng of template DNA, 3  $\mu$ g of dsRNA YFP;
- Lane 6: 50 ng of template DNA, 5  $\mu$ g of dsRNA YFP;
- 15 Lane 7: without template DNA, 1.5 dsRNA YFP;
- Lane 8: 50 ng of template DNA, without dsRNA;
- Lane 9: 50 ng of template DNA, 0.5  $\mu$ g of dsRNA CMV5;
- Lane 10: 50 ng of template DNA, 1.5  $\mu$ g of dsRNA CMV5;
- Lane 11: 50 ng of template DNA, 3  $\mu$ g of dsRNA CMV5;
- 20 Lane 12: 50 ng of template DNA, 5  $\mu$ g of dsRNA CMV5;

It emerged that the amount of transcript was reduced markedly in the presence of dsRNA with sequence homology in comparison with the control reaction without dsRNA and with the reactions with dsRNA YFP without sequence homology. The positive control in lane 2 shows that radioactive transcript was formed upon the *in-vitro* transcription with HeLa nuclear extract. The reaction is used for comparison with the transcription reactions which had been incubated in the presence of dsRNA. Lanes 3 to 6 show that the addition of non-sequentially-specific dsRNA YFP had no effect on the amount of transcript formed. Lanes 9 to 12 show that the addition of an amount of between 1.5 and 3  $\mu$ g of sequentially-specific dsRNA CMV5 leads to a reduction in the amount of transcript formed. In order to exclude that the effects observed are based not on the dsRNA but on any contamination which might have been carried along accidentally during the preparation of the dsRNA,

a further control was carried out. Single-stranded RNA was transcribed as described above and subsequently subjected to the RNase treatment. It was demonstrated by means of native polyacrylamide gel electrophoresis that the ssRNA had been degraded completely. This reaction was subjected to phenol extraction and ethanol precipitation and subsequently taken up in PE buffer, as were the hybridization reactions. This gave a sample which contained no RNA but had been treated with the same enzymes and buffers as the dsRNA. Lane 8 shows that the addition of this sample had no effect on transcription. The reduction of the transcript upon addition of sequence-specific dsRNA can therefore be ascribed unequivocally to the dsRNA itself. The reduction of the amount of transcript of a gene in the presence of dsRNA in a human transcription system indicates an inhibition of the expression of the gene in question. This effect can be attributed to a novel mechanism caused by the dsRNA.

Use example 2:

The test system used for these *in-vivo* experiments was the murine fibroblast cell line NIH3T3, ATCC CRL-1658. The YFP gene was introduced into the nuclei with the aid of microinjection. Expression of YFP was studied under the effect of simultaneously cotransfected dsRNA with sequence homology. This dsRNA YFP shows homology with the 5'-region of the YFP gene over a length of 315 bp. The nucleotide sequence of a strand of the dsRNA YFP is shown in sequence listing No. 5. Evaluation under the fluorescence microscope was carried out 3 hours after injection with reference to the greenish-yellow fluorescence of the YFP formed.

Construction of the template plasmid, and preparation of the dsRNA:

A plasmid was constructed following the same principle as described in use example 1 to act as template for the production of the YFP dsRNA by means of T7 and SP6

in-vitro transcription. Using the primer *Eco\_T7\_YFP* as shown in sequence listing No. 6 and *Bam\_SP6\_YFP* as shown in sequence listing No. 7, the desired gene fragment was amplified by PCR and used analogously to the above description for preparing the dsRNA. The dsRNA YFP obtained is identical to the dsRNA used in use example 1 as non-sequence-specific control.

A dsRNA linked chemically at the 3' end of the RNA as shown in sequence listing No. 8 to the 5' end of the complementary RNA via a C18 linker group was prepared (L-dsRNA). To this end, synthons modified by disulfide bridges were used. The 3'-terminal synthon is bound to the solid support via the 3' carbon with an aliphatic linker group via a disulfide bridge. In the 5'-terminal synthon of the complementary oligoribonucleotide which is complementary to the 3'-terminal synthon of the one oligoribonucleotide, the 5'-trityl protecting group is bound via a further aliphatic linker and a disulfide bridge. Following synthesis of the two single strands, removal of the protecting groups and hybridization of the complementary oligoribonucleotides, the thiol groups which form are brought into spatial vicinity. The single strands are linked to each other by oxidation via their aliphatic linkers and a disulfide bridge. This is followed by purification with the aid of HPLC.

#### Preparation of the cell cultures:

The cells were incubated in DMEM supplemented with 4.5 g/l glucose, 10% fetal bovine serum in culture dishes at 37°C under a 7.5% CO<sub>2</sub> atmosphere and passaged before reaching confluence. The cells were detached with trypsin/EDTA. To prepare for microinjection, the cells were transferred into Petri dishes and incubated further until microcolonies formed.

Microinjection:

For the microinjection, the culture dishes were removed from the incubator for approximately 10 minutes. Approximately 50 nuclei were injected singly per  
5 reaction within a marked area using the AIS microinjection system from Carl Zeiss, Göttingen, Germany. The cells were subsequently incubated for three more hours. For the microinjection, borosilicate glass capillaries from Hilgenberg GmbH, Malsfeld,  
10 Germany, with a diameter of less than 0.5  $\mu\text{m}$  at the tip were prepared. The microinjection was carried out using a micromanipulator from Narishige Scientific Instrument Lab., Tokyo, Japan. The injection time was 0.8 seconds and the pressure was approximately 100 hPa. The  
15 transfection was carried out using the plasmid pCDNA YFP, which contains an approximately 800 bp *Bam*HI/*Eco*RI fragment with the YFP gene in vector pCDNA3. The samples injected into the nuclei contained 0.01  $\mu\text{g}/\mu\text{l}$  of pCDNA-YFP and Texas Red coupled to dextran-70000 in  
20 14 mM NaCl, 3 mM KCl, 10 mM KPO<sub>4</sub> [sic], pH 7.5. Approximately 100 pl of RNA with a concentration of 1  $\mu\text{M}$  or, in the case of the L-dsRNA, 375  $\mu\text{M}$  were additionally added.

25 The cells were studied under a fluorescence microscope with excitation with the light of the excitation wavelength of Texas Red, 568 nm, or of YFP, 488 nm. Individual cells were documented by means of a digital camers. Figures 4a-e show the result for NIH3T3 cells.  
30 In the cells shown in Fig. 4a, sense-YFP-ssRNA has been injected, in Fig. 4b antisense-YFP-ssRNA, in Fig. 4c dsRNA-YFP, in Fig. 4d no RNA and in Fig. 4e L-dsRNA.

The field on the left shows in each case the  
35 fluorescence of cells with excitation at 568 nm. The fluorescence of the same cells at an excitation of 488 nm is seen on the right. The Texas Red fluorescence of all the cells shown demonstrates that the injection solution had been applied successfully into the nuclei

- 5 The right fields of each of figures 4a and 4b show that YFP expression was not visibly inhibited when the single-stranded RNA was injected into the nuclei. The right field of Fig. 4c shows cells whose YFP fluorescence was no longer detectable after the  
10 injection of dsRNA-YFP. Fig. 4d shows cells into which no RNA had been injected, as control. The cell shown in fig. 4e shows YFP fluorescence which can no longer be detected owing to the injection of the L-dsRNA which shows regions with sequence homology to the YFP gene.  
15 This result demonstrates that even shorter dsRNAs can be used for specifically inhibiting gene expression in mammals when the double strands are stabilized by chemically linking the single strands.

## Literatur:

Asanuma, H., Ito, T., Yoshida, T., Liang, X. & Komiyama, M.  
(1999). Photoregulation der Bildung und Dissoziation ei-  
5 nes DNA-Duplexes durch *cis-trans*-Isomerisierung einer  
Azobenzoleinheit. *Angew. Chem.* **111**, 2547-2549.

Azhayeva, E., Azhayev, A., Auriola, S., Tengvall, U., Urtti,  
A. & Lönnberg, H. (1997). Inhibitory properties of double  
10 helix forming circular oligonucleotides. *Nucl. Acids Res.*  
**25**, 4954-4961.

Castelli, J., Wood, K.A. & Youle, R.J. (1998). The 2-5A system  
in viral infection and apoptosis. *Biomed. Pharmacother.*  
15 **52**, 386-390.

Dolinnaya, N.G., Blumenfeld, M., Merenkova, I., Oretskaya,  
T.S., Krynetskaya, N.F., Ivanovskaya, M.G., Vasseur, M. &  
Shabarova, Z.A. (1993). Oligonucleotide circularization  
20 by template-directed chemical ligation. *Nucl. Acids Res.*  
**21**, 5403-5407.

Expert-Bezancon, A., Milet, M. & Carbon, P. (1983). Precise  
localization of several covalent RNA-RNA cross-link in  
25 *Escherichia coli* 16S RNA. *Eur. J. Biochem.* **136**, 267-274.

Fire, A., Xu, S., Montgomery, M.K., Kostas, S.A., Driver, S.E.  
& Mello, C.C. (1998). Potent and specific genetic inter-  
ference by double-stranded RNA in *Caenorhabditis elegans*.  
30 *Nature* **391**, 806-811.

Gao, H., Yang, M., Patel, R. & Cook, A.F. (1995). Circulaiza-  
tion of oligonucleotides by disulfide bridge formation.  
*Nucl. Acids Res.* **23**, 2025-2029.

5 Gryaznov, S.M. & Letsinger, R.L. (1993). Template controlled  
coupling and recombination of oligonucleotide blocks con-  
taining thiophosphoryl groups. *Nucl. Acids Res.* **21**, 1403-  
1408.

10 Kaufman, R.J. (1999). Double-stranded RNA-activated protein  
kinase mediates virus-induced apoptosis: A new role for  
an old actor. *Proc. Natl. Acad. Sci. USA* **96**, 11693-11695.

15 Lipson, S.E. & Hearst, J.E. (1988). Psoralen cross-linking of  
ribosomal RNA. In *Methods in Enzymology* Anonymous pp.  
330-341.

20 Liu, Z.R., Sargueil, B. & Smith, C.W. (1998). Detection of a  
novel ATP-dependent cross-linked protein at the 5' splice  
site-U1 small nuclear RNA duplex by methylene blue-  
mediated photo-cross-linking. *Mol. Cell. Biol.* **18**, 6910-  
6920.

25 Micura, R. (1999). Cyclic oligoribonucleotides (RNA) by solid-  
phase synthesis. *Chem. Eur. J.* **5**, 2077-2082.

30 Skripkin, E., Isel, C., Marquet, R., Ehresmann, B. & Ehres-  
mann, C. (1996). Psoralen crosslinking between human im-  
munodeficiency virus type 1 RNA and primer tRNA<sub>3</sub><sup>Lys</sup>. *Nucl.*  
*Acids Res.* **24**, 509-514.

- Wang, S. & Kool, E.T. (1994). Circular RNA oligonucleotides. Synthesis, nucleic acid binding properties, and a comparison with circular DNAs. *Nucl. Acids Res.* **22**, 2326-2333.
- 5 Wang, Z. & Rana, T.M. (1996). RNA conformation in the Tat-TAR complex determined by site-specific photo-cross-linking. *Biochem.* **35**, 6491-6499.
- Watkins, K.P. & Agabian, N. (1991). In vivo UV cross-linking  
10 of U snRNAs that participate in trypanosome trans-splicing. *Genes & Development* **5**, 1859-1869.
- Wengel, J. (1999). Synthesis of 3'-C- and 4'-C-branched oligodeoxynucleotides and the development of locked nucleic  
15 acid (LNA). *Acc. Chem. Res.* **32**, 301-310.
- Zwieb, C., Ross, A., Rinke, J., Meinke, M. & Brimacombe, R. (1978). Evidence for RNA-RNA cross-link formation in  
20 *Escherichia coli* ribosomes. *Nucl. Acids Res.* **5**, 2705-2720.